


Lumiracoxib

Chemical Properties

CAS No. :	220991-20-8
Formula:	C ₁₅ H ₁₃ ClFNO ₂
Molecular Weight:	293.72
Appearance:	no data available
Storage:	store at low temperature
	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Lumiracoxib (Prexige) is a novel, selective COX-2 inhibitor with IC ₅₀ and K _i of 0.14 μM and 0.06 μM, exhibits 515-fold selectivity over COX-1. Phase 4.
Targets(IC ₅₀)	COX
In vitro	Lumiracoxib has an IC ₅₀ of 0.14 μM in COX-2-expressing dermal fibroblasts, but caused no inhibition of COX-1 at concentrations up to 30 μM (HEK 293 cells transfected with human COX-1). In a human whole blood assay, IC ₅₀ values for Lumiracoxib are 0.13 μM for COX-2 and 67 μM for COX-1 (COX-1/COX-2 selectivity ratio 515).
In vivo	Lumiracoxib is a highly selective COX-2 inhibitor with anti-inflammatory, analgesic and antipyretic activities comparable with diclofenac, the reference NSAID, but with much improved gastrointestinal safety. Lumiracoxib is rapidly absorbed following oral administration in rats with peak plasma levels being reached between 0.5 and 1 h. Efficacy of Lumiracoxib in rat models of hyperalgesia, oedema, pyresis and arthritis is dose-dependent and similar to diclofenac. However, consistent with its low COX-1 inhibitory activity, Lumiracoxib at a dose of 100 mg/kg orally causes no ulcers and is significantly less ulcerogenic than diclofenac.
Animal Research	Animal Models: Female Lewis rats Formulation: Sterile phosphate-buffered saline Dosages: 0.2–2 mg/kg Administration: Oral gavage

Solubility Information

Solubility	H ₂ O: < 1 mg/mL (insoluble or slightly soluble), Ethanol: < 1 mg/mL (insoluble or slightly soluble), DMSO: 55 mg/mL (187.25 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.4046 mL	17.023 mL	34.046 mL
5 mM	0.6809 mL	3.4046 mL	6.8092 mL
10 mM	0.3405 mL	1.7023 mL	3.4046 mL
50 mM	0.0681 mL	0.3405 mL	0.6809 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Esser R, et al. Br J Pharmacol, 2005, 144(4), 538-550.

Sun J, Zhang L, Zhang L, et al. A validated UHPLC-MS/MS method for simultaneous determination of lumiracoxib and its hydroxylation and acyl glucuronidation metabolites in rat plasma: Application to a pharmacokinetic study. Journal of Pharmaceutical and Biomedical Analysis. 2021, 201: 114105.

Sun J, Zhang L, Zhang L, et al. A validated UHPLC-MS/MS method for simultaneous determination of lumiracoxib and its hydroxylation and acyl glucuronidation metabolites in rat plasma: application to a pharmacokinetic study. Journal of Pharmaceutical and Biomedical Analysis. 2021: 114105.

Jiao W, Zhao X, Wu G, et al. Bioactivation of Lumiracoxib in Human Liver Microsomes: Formation of GSH-and Amino Adducts Through Acyl Glucuronide[J]. Drug Testing and Analysis. 2020, 12(6): 827-835.

Sun J, Zhang L, Zhang L, et al. A validated UHPLC-MS/MS method for simultaneous determination of lumiracoxib and its hydroxylation and acyl glucuronidation metabolites in rat plasma: application to a pharmacokinetic study [J]. Journal of Pharmaceutical and Biomedical Analysis. 2021: 114105.

Jiao W, Zhao X, Wu G, et al. Bioactivation of Lumiracoxib in Human Liver Microsomes: Formation of GSH-and Amino Adducts Through Acyl Glucuronide. Drug Testing and Analysis. 2020, 12(6): 827-835.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481