Data Sheet (Cat.No.TMPH-02017)



Reelin Protein, Human, Recombinant (E. coli, His)

General Information

Synonyms: RELN;Reelin

Protein Construction: 26-254 aa

Species: Human

Expression Host: E. coli

Accession: P78509

Molecular Weight: 28.9 kDa (predicted)

AAGYYPRFSPFFLCTHHGELEGDGEQGEVLISLHIAGNPTYYVPGQEYHVTISTSTFFDGLLVTGLYTSTSVQA

AA Sequence: SQSIGGSSAFGFGIMSDHQFGNQFMCSVVASHVSHLPTTNLSFIWIAPPAGTGCVNFMATATHRGQVIFKDA

LAQQLCEQGAPTDVTVHPHLAEIHSDSIILRDDFDSYHQLQLNPNIWVECNNCETGEQCGAIMHGNAVTFCE PYGPRELITT

QC Testing

Biological Activity: Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you

require protein activity, we recommend choosing the eukaryotic expression version first.

Purity: > 90% as determined by SDS-PAGE.

Endotoxin: $< 1.0 \text{ EU/}\mu\text{g}$ of the protein as determined by the LAL method.

Formulation: Tris-based buffer, 50% glycerol

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Shipping:

In general, Lyophilized powders are shipping with blue ice. Solutions are shipping with dry ice.

Protein Background

Extracellular matrix serine protease that plays a role in layering of neurons in the cerebral cortex and cerebellum. Regulates microtubule function in neurons and neuronal migration. Affects migration of sympathetic preganglionic neurons in the spinal cord, where it seems to act as a barrier to neuronal migration. Enzymatic activity is important for the modulation of cell adhesion. Binding to the extracellular domains of lipoprotein

Page 1 of 2 www.targetmol.com

receptors VLDLR and LRP8/APOER2 induces tyrosine phosphorylation of DAB1 and modulation of TAU phosphorylation.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481

Page 2 of 2 www.targetmol.com