

Complement C3 Protein, Mouse, Recombinant (His & SUMO)

General Information

Synonyms: C3;HSE-MSF;Complement C3

Protein Construction: 1321-1663 aa

Species: Mouse

Expression Host: E. coli

Accession: P01027

Molecular Weight: 55.3 kDa (predicted)

SEETKQNEAFSLTAKGKGRGTLVVAVYHAKLKSKVTCKKFDLRSIRPAPETAKPPEAKNTMFLEICTKYLGDVDATMSILDISMMTGFAPDTKDLLELASGVDRYISKYEMNKAFSNKNTLIIYLEKISHTEDCLTFKVHQYFNVGLIQPGSVKVYSYYNLEESCTRFYHPEKDDGMLSCLCHSEMCRAEENCFMQSQEKinLNVRLDKACEPGVDYVYKTELNIELLDDFDEYTMТИQQVIKSGSDEVQAGQQRKFISHIKCRNALKLQKGKKYLMWGLSSDWGEPNKPNTSYIIGKDTWVEHWPEAECQDQKYQKCEELGAFTESMVVYGCN

QC Testing

Biological Activity: Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.

Purity: > 90% as determined by SDS-PAGE.

Endotoxin: < 1.0 EU/μg of the protein as determined by the LAL method.

Formulation: Tris-based buffer, 50% glycerol

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Shipping:

In general, Lyophilized powders are shipping with blue ice. Solutions are shipping with dry ice.

Protein Background

C3 plays a central role in the activation of the complement system. Its processing by C3 convertase is the central reaction in both classical and alternative complement pathways. After activation C3b can bind covalently, via its reactive thioester, to cell surface carbohydrates or immune aggregates.; Derived from proteolytic degradation of

complement C3, C3a anaphylatoxin is a mediator of local inflammatory process. In chronic inflammation, acts as a chemoattractant for neutrophils. It induces the contraction of smooth muscle, increases vascular permeability and causes histamine release from mast cells and basophilic leukocytes. The short isoform has B-cell stimulatory activity.; Acts as a chemoattractant for neutrophils in chronic inflammation.; adipogenic hormone that stimulates triglyceride (TG) synthesis and glucose transport in adipocytes, regulating fat storage and playing a role in postprandial TG clearance. Appears to stimulate TG synthesis via activation of the PLC, MAPK and AKT signaling pathways. Ligand for C5AR2. Promotes the phosphorylation, ARRB2-mediated internalization and recycling of C5AR2.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481