

## OX40/TNFRSF4 Protein, Mouse, Recombinant (His)

### General Information

Synonyms: Txgp1;OX40;Tnfrsf4;Tumor necrosis factor receptor superfamily member 4;CD134

Protein Construction: Val20-Pro211

Species: Mouse

Expression Host: HEK293 Cells

Accession: P47741

Molecular Weight: 40 KDa (reducing condition)

AA Sequence: Val20-Pro211

### QC Testing

Biological Activity: Loaded Mouse OX40L-His on AR2G Biosensor, can bind Mouse OX40-His with an affinity constant of 0.16  $\mu$ M as determined in BLI assay. (Regularly tested)

Purity: Greater than 95% as determined by reducing SDS-PAGE. (QC verified)

Endotoxin: < 0.1 ng/ $\mu$ g (1 EU/ $\mu$ g) as determined by LAL test.

Formulation: Lyophilized from a solution filtered through a 0.22  $\mu$ m filter, containing PBS, pH 7.4.

### Preparation and Storage

#### Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100  $\mu$ g/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

#### Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

#### Shipping:

In general, Lyophilized powders are shipping with blue ice. Solutions are shipping with dry ice.

### Protein Background

OX40, also termed CD134 and TNFRSF4, is a T cell co-stimulatory molecule of the TNF receptor superfamily which plays a key role in the survival and homeostasis of effector and memory T cells. OX40 is expressed on CD4+ and CD8+ T cells upon engagement of the TCR by antigen presenting cells along with co-stimulation by CD40-CD40 Ligand and CD28-B7. The interaction between OX40 and OX40 ligand (OX40L) will occur when activated T cells bind to professional antigen-presenting cells (APCs). The T-cell functions, including cytokine production, expansion, and survival, are then enhanced by the OX40 costimulatory signals. OX40 signals are critical for

controlling the function and differentiation of Foxp3+ regulatory T cells. OX40-OX40L interaction regulates T-cell tolerance, peripheral T-cell homeostasis, and T-cell-mediated inflammatory diseases.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481