

OBhFc1 Protein, Human, Recombinant (His)

General Information

Synonyms:	Suppressor of Cdc Thirteen Homolog;CST Complex Subunit STN1; Oligonucleotide/Oligosaccharide-Binding Fold-Containing Protein 1;OBFC1;STN1
Protein Construction:	Met1-Phe368
Species:	Human
Expression Host:	<i>E. coli</i>
Accession:	AAH17400.1
Molecular Weight:	44 KDa (reducing condition)
AA Sequence:	Met1-Phe368

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 80% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing 20 mM Tris-HCl, 1 mM DTT, 5% Trehalose, pH 8.0.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Shipping:

In general, Lyophilized powders are shipping with blue ice. Solutions are shipping with dry ice.

Protein Background

CST Complex Subunit STN1 (OBFC1) is a 368 amino acid protein that contains one OB DNA-binding domain. It is a member of the STN1 family. OBFC1 is component of the CST complex, a complex that binds to single-stranded DNA and is required to protect telomeres from DNA degradation. The CST complex binds single-stranded DNA with high affinity in a sequence-independent manner, while isolated subunits bind DNA with low affinity by themselves. In addition to telomere protection, the CST complex has probably a more general role in DNA metabolism at non-

telomeric sites.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481