

## CD177 Protein, Human, Recombinant (His)

### General Information

Synonyms:	PRV-1;HNA-2a;Human Neutrophil Alloantigen 2a;CD177;Polycythemia Rubra Vera Protein 1; NB1;NB1 Glycoprotein;NB1 GP;CD177 Antigen;PRV1
Protein Construction:	Leu22-Gly407
Species:	Human
Expression Host:	HEK293 Cells
Accession:	AAH29167.1
Molecular Weight:	55 KDa (reducing condition)
AA Sequence:	Leu22-Gly407

### QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing 20 mM PB, 150 mM NaCl, pH 7.2.

### Preparation and Storage

#### Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

#### Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

#### Shipping:

In general, Lyophilized powders are shipping with blue ice. Solutions are shipping with dry ice.

### Protein Background

CD177 is polymorphic and has at least two alleles: PRV1 and NB1. Human PRV1 is a Glycosyl-Phosphatidylinositol (GPI)-linked cell surface glycoprotein that belongs to the uPAR/CD59/Ly6 family of receptors. PRV1 is expressed by neutrophils and neutrophil precursors, and changes in expression serve as diagnostic markers for myeloproliferative disorders such as polycythemia vera and essential thrombocythemia. PRV1 may also be expressed by Erythroblasts, B cells, and Monocytes. NB1, a Glycosyl-Phosphatidylinositol (GPI)-linked cell surface

glycoprotein, was first described in a case of neonatal alloimmune neutropenia. It is reported that CD177 functions as a novel heterophilic binding partner that engages PECAM-1 in membrane-proximal IgD6.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481