

HMGB2 Protein, Human, Recombinant (His)

General Information

| | |
|-----------------------|---|
| Synonyms: | High Mobility Group Protein B2;HMG2;High Mobility Group Protein 2;HMG-2;HMGB2 |
| Protein Construction: | Gly2-Glu209 |
| Species: | Human |
| Expression Host: | HEK293 Cells |
| Accession: | P26583 |
| Molecular Weight: | 28 KDa (reducing condition) |
| AA Sequence: | Gly2-Glu209 |

QC Testing

| | |
|----------------------|---|
| Biological Activity: | Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first. |
| Purity: | Greater than 95% as determined by reducing SDS-PAGE. (QC verified) |
| Endotoxin: | < 0.1 ng/µg (1 EU/µg) as determined by LAL test. |
| Formulation: | Lyophilized from a solution filtered through a 0.22 µm filter, containing 20 mM Tris-HCl, 100 mM NaCl, pH 8.0. |

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 µg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Shipping:

In general, Lyophilized powders are shipping with blue ice. Solutions are shipping with dry ice.

Protein Background

High Mobility Group Protein B2 (HMGB2) belongs to the non-histone chromosomal high-mobility group protein family. Members of this family are chromatin-associated and widely spread in the nucleus of higher eukaryotic cells. HMGB2 contains 2 HMG box DNA-binding domains. It is associated with chromatin and has the ability to bend DNA, preferentially single-stranded DNA. It is shown that HMGB2 is able to efficiently bend DNA and form DNA circles. In addition, HMGB2 is involved in the final ligation step in DNA end-joining processes of DNA double-

strand breaks repair and V(D)J recombination.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481