

EIF4EBP2 Protein, Human, Recombinant (His)

General Information

| | |
|-----------------------|--|
| Synonyms: | Eukaryotic Translation Initiation Factor 4E-Binding Protein 2; EIF4EBP2; eIF4E-Binding Protein 2; 4E-BP2 |
| Protein Construction: | Met1-Ile120 |
| Species: | Human |
| Expression Host: | E. coli |
| Accession: | Q13542 |
| Molecular Weight: | 17 KDa (reducing condition) |
| AA Sequence: | Met1-Ile120 |

QC Testing

| | |
|----------------------|---|
| Biological Activity: | Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first. |
| Purity: | Greater than 90% as determined by reducing SDS-PAGE. (QC verified) |
| Endotoxin: | < 0.1 ng/μg (1 EU/μg) as determined by LAL test. |
| Formulation: | Lyophilized from a solution filtered through a 0.22 μm filter, containing 20 mM Tris-HCl, 150 mM NaCl, pH 8.0 . |

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Shipping:

In general, Lyophilized powders are shipping with blue ice. Solutions are shipping with dry ice.

Protein Background

Eukaryotic Translation Initiation Factor 4E-Binding Protein 2 (EIF4EBP2) is a member of the Eukaryotic Translation Initiation Factor 4E Binding Protein Family. EIF4EBP2 regulates eIF4E activity by preventing its assembly into the eIF4F complex, mediates the regulation of protein translation by hormones, growth factors and other stimuli that signal through the MAP kinase pathway. This regulation of is associated to cell proliferation, cell differentiation and viral infection. Phosphorylated EIF4EBP2 on serine and threonine residues in response to insulin, EGF and

PDGF.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481