

NKG2D/CD314 Protein, Human, Recombinant (hFc & Flag), FITC-Labeled

General Information

Synonyms:	NKG2D;KLR;CD314;D12S2489E;NKG2-D
Protein Construction:	Phe78-Val216
Species:	Human
Expression Host:	HEK293 Cells
Accession:	P26718
Molecular Weight:	43.4 kDa (predicted). Due to glycosylation, the protein migrates to 50-70 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
Endotoxin:	< 1 EU/μg by the LAL method.
Formulation:	Supplied as 0.22 μm filtered solution in PBS (pH 7.4).

Preparation and Storage

Stability & Storage:	It is recommended to store the product under sterile conditions at -70°C or lower. Samples are stable for up to 12 months at -80°C. Please avoid multiple freeze-thaw cycles and store products in aliquots.
Shipping:	Shipping with blue ice.

Protein Background

NKG2D is a type II transmembrane glycoprotein having an extracellular lectin-like domain. This domain lacks the recognizable calcium-binding sites found in true C-type lectins and binds protein rather than carbohydrate ligands. Human NKG2D is expressed on CD8 alpha beta T cells, gamma δ T cells, NK cells and NKT cells.

Reference

Spear P, et al. NKG2D CAR T-cell therapy inhibits the growth of NKG2D ligand heterogeneous tumors[J]. Immunology and Cell Biology, 2013, 91(6):435-440.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481