

SIRP alpha V8 Protein, Human, Recombinant (His & Avi), Biotinylated

General Information

Synonyms:	P84;MFR;MYD1;SHPS-1;SIRP alpha;SIRP α ;SIRP α ;MYD-1;PTPNS1;SIRP α V8;SHPS1;CD172a;BIT;SIRPA
Protein Construction:	Glu31-Arg369(S44L, S50T, I52T, H54R, V57A)
Species:	Human
Expression Host:	HEK293 Cells
Accession:	ATD50864.1
Molecular Weight:	40.1 kDa (predicted). Due to glycosylation, the protein migrates to 52-68 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	Immobilized Human CD47, hFc Tag at 1 μ g/ml (100 μ l/well) on the plate. Dose response curve for Biotinylated Human SIRP alpha V8, His Tag with the EC50 of 0.53 μ g/ml determined by ELISA.
Purity:	> 95% as determined by Tris-Bis PAGE; > 90% as determined by HPLC
Endotoxin:	< 1 EU/ μ g by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μ m filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μ g/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Shipping:

In general, Lyophilized powders are shipping with blue ice.

Protein Background

Signal regulatory protein α (SIRP α) is a regulatory membrane glycoprotein from SIRP family expressed mainly by myeloid cells and also by stem cells or neurons. SIRP α acts as inhibitory receptor and interacts with a broadly expressed transmembrane protein CD47 also called the "don't eat me" signal. Cancer cells highly expressed CD47 that activate SIRP α and inhibit macrophage-mediated destruction.

Reference

Weiskopf K, et al. Engineered SIRP α variants as immunotherapeutic adjuvants to anticancer antibodies[J]. Science, 2013, 341(6141):88-91.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481