

## SFRP2 Protein, Cynomolgus, Recombinant (His)

### General Information

Synonyms:	Sdf5;SARP-1;sFRP-2;Sarp1;Sfrp2
Protein Construction:	Leu25-Cys295
Species:	Cynomolgus
Expression Host:	HEK293 Cells
Accession:	G8F2Z7
Molecular Weight:	32.16 kDa (predicted). Due to glycosylation, the protein migrates to 33-38 kDa based on Tris-Bis PAGE result.

### QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 95% as determined by Tris-Bis PAGE
Endotoxin:	< 1 EU/μg by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

### Preparation and Storage

Reconstitution:	Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.
Stability & Storage:	It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.
Shipping:	In general, Lyophilized powders are shipping with blue ice.

### Protein Background

As biomarkers, DNA methylation is used to detect colorectal cancer (CRC) and make assessment of CRC prognosis. The published findings showed the association between the methylation of SFRP1, SFRP2, and WIF1, located in the Wnt signaling pathway, and the prognosis of CRC were not consistent. SFRP1, SFRP2, and WIF1 were frequently hypermethylated in CRC tumor tissues. It was apparent that the promoter hypermethylation of SFRP2 and co-hypermethylation of SFRP1 and SFRP2 might be considered as independent prognostic predictors for survival

advantage of postoperative CRC patients.

Reference

Liu X, et al. DNA methylation of SFRP1, SFRP2, and WIF1 and prognosis of postoperative colorectal cancer patients. BMC Cancer. 2019 Dec 12;19(1):1212. doi: 10.1186/s12885-019-6436-0. PMID: 31830937; PMCID: PMC6909551.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286    E\_mail: [info@targetmol.com](mailto:info@targetmol.com)    Address: 34 Washington Street, Wellesley Hills, MA 02481