

## SLAMF6 Protein, Mouse, Recombinant (aa 31-239, His)

## General Information

Synonyms:	Ly108; ANK-T-B-antigen; NTB-A; KALI; NK-T-B-antigen; KAL1b; CD352; SLAMF6; SF2000
Protein Construction:	Glu31-Asn239
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	Q9ET39-1
Molecular Weight:	24.03 kDa (predicted). Due to glycosylation, the protein migrates to 40-55 kDa based on Tris-Bis PAGE result.

## QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
Endotoxin:	< 1 EU/μg by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

## Preparation and Storage

## Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

## Stability &amp; Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

## Shipping:

In general, Lyophilized powders are shipping with blue ice.

## Protein Background

SLAMF6 (signaling lymphocyte activation molecule 6) (Ly108 in mice, NTB-A or SF2000 in humans) is a homophilic receptor belonging to the superfamily immunoglobulin (Ig) domain-containing molecules. It is known to be widely and exclusively expressed on hematopoietic cells. The SLAMF6 intracellular portion is characterized by two ITSMs that act as binding sites for adaptor molecules such as SAP and EAT-2.

### Reference

Dragovich MA, et al. SLAMF6 clustering is required to augment T cell activation. PLoS One. 2019 Jun 14;14(6): e0218109. doi: 10.1371/journal.pone.0218109. PMID: 31199820; PMCID: PMC6568412.

Yigit B, et al. SLAMF6 as a Regulator of Exhausted CD8+ T Cells in Cancer. Cancer Immunol Res. 2019 Sep;7(9): 1485-1496. doi: 10.1158/2326-6066.CIR-18-0664. Epub 2019 Jul 17. PMID: 31315913.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286    E\_mail: info@targetmol.com    Address: 34 Washington Street, Wellesley Hills, MA 02481