

GPVI Protein, Human, Recombinant (His)

General Information

Synonyms: Glycoprotein 6;Gp6;MGC138168;GPVI

Protein Construction: Gln21-Lys267

Species: Human

Expression Host: HEK293 Cells

Accession: Q9HCN6-1

Molecular Weight: 28 kDa (predicted). Due to glycosylation, the protein migrates to 50-60 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity: Immobilized Human GPVI, His Tag at 0.5 μ g/ml (100 μ l/Well) on the plate. Dose response curve for Anti-GPVI Antibody, hFc Tag with the EC50 of 4.2ng/ml determined by ELISA.

Purity: > 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC

Endotoxin: < 1 EU/ μ g by the LAL method.

Formulation: Lyophilized from a solution filtered through a 0.22 μ m filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μ g/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Shipping:

In general, Lyophilized powders are shipping with blue ice.

Protein Background

Although platelets are best known for their role in hemostasis, they are also crucial in development, host defense, inflammation, and tissue repair. Many of these roles are regulated by the immune-like receptors glycoprotein VI (GPVI) and C-type lectin receptor 2 (CLEC-2), which signal through an immunoreceptor tyrosine-based activation motif (ITAM). GPVI is activated by collagen in the subendothelial matrix, by fibrin and fibrinogen in the thrombus, and by a remarkable number of other ligands.

Reference

Rayes J, et al. Functional significance of the platelet immune receptors GPVI and CLEC-2. *J Clin Invest.* 2019 Jan 2; 129(1):12-23. doi: 10.1172/JCI122955. Epub 2019 Jan 2. PMID: 30601137; PMCID: PMC6307936.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481