

Transferrin Receptor/TFRC Protein, Cynomolgus, Recombinant (His)

General Information

Synonyms: TFR;T9;CD71;TfR1;sTfR;TFRC;TR;p90

Protein Construction: Cys89-Phe760

Species: Cynomolgus

Expression Host: HEK293 Cells

Accession: XP_005545315.1

Molecular Weight: 76.3 kDa (predicted). Due to glycosylation, the protein migrates to 78-80 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity: Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.

Purity: > 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC

Endotoxin: < 1 EU/μg by the LAL method.

Formulation: Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Shipping:

In general, Lyophilized powders are shipping with blue ice.

Protein Background

The transferrin receptor (CD71) is known as a receptor for IgA1 on mesangial cells. The transcript and protein expression of glomerular CD71 was determined by real-time polymerase chain reaction and immunohistochemistry. Human mesangial cells (HMCs) were treated with sera from IgAN patients and expression levels of CD71 and inflammatory cytokine markers were compared according to disease status.

Reference

Jhee JH, et al. CD71 mesangial IgA1 receptor and the progression of IgA nephropathy. *Transl Res*. 2021 Apr;230:34-43. doi: 10.1016/j.trsl.2020.10.007. Epub 2020 Oct 24. PMID: 33122053.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481