

IL-22 Protein, Human, Recombinant (HEK293)

General Information

Synonyms: IL-D110; interleukin 22; IL22; IL-22; TIFa; ILTIF; IL-TIF; TIFIL-23; IL-21; zcyto18

Protein Construction: Ala34-Ile179

Species: Human

Expression Host: HEK293 Cells

Accession: Q9GZX6

Molecular Weight: 16~28 kDa (Reducing conditions)

QC Testing

Biological Activity: Measured by its ability to induce IL-10 secretion in COLO 205 (human colon carcinoma cells). The ED 50 for this effect is less than 0.3 ng/ml.

Purity: > 95% as determined by SDS-PAGE

Endotoxin: < 0.2 EU/μg of protein as determined by the LAL method.

Formulation: Lyophilized from a 0.2 μm filtered solution in PBS.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Shipping:

In general, Lyophilized powders are shipping with blue ice. Solutions are shipping with dry ice.

Protein Background

Interleukin-22(IL-22) belongs to a group of cytokines called the IL-10 family or IL-10 superfamily (including IL-19, IL-20, IL-24, and IL-26) which are a class of potent mediators of cellular inflammatory responses. It shares use of IL-10R2 in cell signaling with other members of this family, such as IL-10, IL-26, IL-28A/B and IL-29. IL-22 is produced by activated DC and T cells and initiates innate immune responses against bacterial pathogens in epithelial cells such as those in the lung and gut. IL-22 along with IL-17 is produced by splenic LTI-like cells and Th17 cells and likely plays a role in the coordinated response of both adaptive and innate immune systems. IL-22 signals through a receptor system consisting of IL-10R-β/CRF2-4 and IL-22R, both of which are members of the

class II cytokine-receptor family.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481