

IL-22 Protein, Mouse, Recombinant (CHO)

General Information

Synonyms: IL-D110;IL-22;IL-21;IL-TIF;ILTIF;TIFIL-23;zcyto18;IL22;TIFa;interleukin 22

Protein Construction: Leu34-Val179

Species: Mouse

Expression Host: CHO Cells

Accession: Q9JJY9

Molecular Weight: 20~30 kDa (Reducing conditions)

QC Testing

Biological Activity: ED 50 < 0.5 ng/ml, measured by its ability to induce IL-10 secretion in COLO 205 (human colon carcinoma cells).

Purity: > 95% as determined by SDS-PAGE; > 95% as determined by HPLC

Endotoxin: < 0.2 EU/μg of protein as determined by the LAL method.

Formulation: Lyophilized from a 0.2 μm filtered solution in PBS.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Shipping:

In general, Lyophilized powders are shipping with blue ice. Solutions are shipping with dry ice.

Protein Background

Interleukin-22 (IL-22) is a member of a group of cytokines called the IL-10 family which include IL-10, IL-19, IL-20, IL-24, and IL-26. IL-22 shares use of the IL-10R2 in cell signaling with other members of this family. IL-22 signals through a receptor system consisting of IL-10R-β/CRF2-4 and IL-22R, both of which are members of the class II cytokine-receptor family. IL-22 is produced by activated DCs and T cells and initiates an innate immune response against bacterial pathogens especially in epithelial cells such as those in the respiratory tract and gut. IL-22 along with IL-17 is rapidly produced by splenic LTi-like cells and can also be produced by Th17 cells, which play a likely role in the coordinated response of both adaptive and innate immune systems.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481