

## FGF-21 Protein, Human, Recombinant

### General Information

Synonyms: FGF21;FGFL;fibroblast growth factor-21

Protein Construction: His29-Ser209

Species: Human

Expression Host: E. coli

Accession: Q9NSA1

Molecular Weight: ~19.5 kDa (Reducing conditions)

### QC Testing

Biological Activity: ED 50 < 0.5 µg/ml, measured by a cell proliferation assay using NIH-3T3 cells in the presence of 1.25 µg/ml mouse Klotho and 10 µg/ml heparin, corresponding to a specific activity of > 2.0 × 10<sup>3</sup> units/mg.

Purity: > 95% as determined by SDS-PAGE; > 95% as determined by HPLC

Endotoxin: < 0.2 EU/µg of protein as determined by the LAL method.

Formulation: Lyophilized from a 0.2 µm filtered solution in PBS.

### Preparation and Storage

#### Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 µg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

#### Stability & Storage:

Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

#### Shipping:

In general, Lyophilized powders are shipping with blue ice. Solutions are shipping with dry ice.

### Protein Background

Fibroblast Growth Factor-21 (FGF-21) is a metabolic cytokine belonging to the heparin-binding FGF family. Along with FGF-19/15 and FGF-23, FGF-21 is categorized as a member of the atypical FGF subfamily, as it must be complexed to the Klotho co-receptor in order to bind to the FGF receptors and activate the downstream signaling pathway. In vivo FGF-21 is expressed in liver, pancreas, adipose tissue, and skeletal muscle, and it plays a central role in the energy metabolism. The expression of FGF-21 is stimulated by free fatty acids and insulin resistant states and is correlated with whole-body insulin resistance. FGF-21 activates glucose uptake in adipocytes and

increases insulin sensitivity, implicating it as a novel target with potential anti-diabetic properties.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481