

ALK-4/ACVR1B Protein, Mouse, Recombinant (hFc)

General Information

Synonyms:	Alk4;ActRIB;Acvrlk4;ActR-IB;6820432J04;SKR2;activin A receptor, type IB
Protein Construction:	A DNA sequence encoding the mouse Acvr1b (NP_031421.1) (Met1-Glu126) was expressed with the Fc region of human IgG1 at the C-terminus. Predicted N terminal: Ser
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	Q3TZF1
Molecular Weight:	38.4 kDa (predicted)

QC Testing

Biological Activity:	Measured by its ability to bind with rat TDGF1-His in a functional ELISA.
Purity:	> 95 % as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Shipping:

In general, Lyophilized powders are shipping with blue ice.

Protein Background

ALK-4 (Activin Receptor-Like Kinase 4) or ACVR1B (Activin A Receptor, type 1B), belongs to the protein kinase superfamily, TKL Ser/Thr protein kinase family, and TGF β receptor subfamily. ALK-4/ACVR1B acts as a transducer of activin or activin like ligands signals. Activin binds to either ACVR2A or ACVR2B and then forms a complex with ACVR1B. The known type II activin receptors include ActRII and ActRIIB, while the main type I activin receptor in mammalian cells is ALK-4 (ActRIB). In the presence of activin, type II and type I receptors form complexes whereby the type II receptors activate ALK-4 through phosphorylation. The activated ALK-4, in turn, transduces signals

downstream by phosphorylation of its effectors, such as Smads, to regulate gene expression and affect cellular phenotype. ALK-4/ACVR1B is an important regulator of vertebrate development, with roles in mesoderm induction, primitive streak formation, gastrulation, dorsoanterior patterning, and left-right axis determination.

Reference

Chen Y, et al. (2005) Developmental analysis of activin-like kinase receptor-4 (ALK4) expression in *Xenopus laevis*. 232(2): 393-8.J.
Massagu. (1998) TGF- SIGNAL TRANSDUCTION. Annual Review of Biochemistry. 67: 753-91.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481