

H Cadherin Protein, Mouse, Recombinant (His)

General Information

Synonyms: Cdht;T-cadherin;cadherin 13;4932416G01Rik;Tcad

Protein Construction: A DNA sequence encoding the mouse Cdh13 (NP_062681.2) (Met1-Ala692) was expressed with a polyhistidine tag at the C-terminus. Predicted N terminal: Asp 23

Species: Mouse

Expression Host: HEK293 Cells

Accession: Q9WTR5

Molecular Weight: 75 kDa (predicted)

QC Testing

Biological Activity: Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.

Purity: > 90 % as determined by SDS-PAGE

Endotoxin: < 1.0 EU/μg of the protein as determined by the LAL method.

Formulation: Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Shipping:

In general, Lyophilized powders are shipping with blue ice.

Protein Background

CDH13, also known as cadherin-13 and H Cadherin, is a member of the cadherin superfamily. CDH13 acts as a negative regulator of axon growth during neural differentiation. It also protects vascular endothelial cells from apoptosis due to oxidative stress, and is associated with resistance to atherosclerosis. CDH13 is localized to the surface of the cell membrane and is anchored by a GPI moiety, rather than by a transmembrane domain. CDH13 gene is hypermethylated in many types of cancer.

Reference

Hart AB. et al., 2012, PLoS One. 7 (8): e42646.
Xu J. et al., 2012, BMC Cancer. 12: 243.
Jo J. et al., 2012, Obesity (Silver Spring). 20 (8): 1683-7.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481