

IL-2RG Protein, Mouse, Recombinant (His & hFc)

General Information

Synonyms:	gamma(c);interleukin 2 receptor, gamma;[g]c;gc;γ(c);CD132;interleukin 2 receptor, γ
Protein Construction:	A DNA sequence encoding the extracellular domain (Met 1-Ala 263) of mouse IL2RG precursor was fused with the C-terminal polyhistidine-tagged Fc region of human IgG1 at the C-terminus. Predicted N terminal: Trp 23
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	P34902
Molecular Weight:	56.3 kDa (predicted); 75-85 kDa (reducing condition, due to glycosylation)

QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 92 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Shipping:

In general, Lyophilized powders are shipping with blue ice.

Protein Background

The common gamma chain (γc) (or CD132), also known as interleukin-2 receptor subunit gamma or IL2RG, is a member of the type I cytokine receptor family expressed on most lymphocyte (white blood cell) populations, and its gene is found on the X-chromosome of mammals. The common gamma chain (γc) (or IL2RG), is a cytokine receptor subunit that is common to the receptor complexes for at least six different interleukin receptors: IL-2, IL-4,

IL-7, IL-9, IL-15, and the interleukin-21 receptor. It is a component of multiple cytokine receptors that are essential for lymphocyte development and function. X-linked severe combined immunodeficiency (X-SCID) is a rare and potentially fatal disease caused by mutations of IL2RG, the gene encoding IL2RG. IL2RG was demonstrated to be a component of the IL-4 receptor based on chemical cross-linking data, the ability of IL2RG to augment IL-4 binding affinity. The observation that IL-2R gamma is a functional component of the IL-4 receptor, together with the finding that IL-2R gamma associates with the IL-7 receptor, begins to elucidate why a deficiency of this common gamma chain (gamma c) has a profound effect on lymphoid function and development, as seen in X-linked severe combined immunodeficiency.

Reference

Russell SM, et al. (1993) Interleukin-2 receptor gamma chain: a functional component of the interleukin-4 receptor. *Science*. 262 (5141): 1880-3.

Miyazaki T, et al. (1994) Functional activation of Jak1 and Jak3 by selective association with IL-2 receptor subunits. *Science*. 266 (5187): 1045-7.

Takeshita T, et al. (1992) Cloning of the gamma chain of the human IL-2 receptor. *Science*. 257 (5068): 379-82.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481