

CD52 Protein, Rat, Recombinant (hFc)

General Information

Synonyms: CD52 molecule

Protein Construction: A DNA sequence encoding the rat ILCD52 (Q63064) (Met1-Ala69) was expressed, fused with the Fc region of human IgG1 at the C-terminus. Predicted N terminal: Gly 24

Species: Rat

Expression Host: HEK293 Cells

Accession: Q63064

Molecular Weight: 31.4 kDa (predicted); 53 kDa (reducing conditions)

QC Testing

Biological Activity: Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.

Purity: > 70 % as determined by SDS-PAGE

Endotoxin: < 1.0 EU/μg of the protein as determined by the LAL method.

Formulation: Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Shipping:

In general, Lyophilized powders are shipping with blue ice.

Protein Background

CD52 / CDW52 is a small glycosylphosphatidylinositol (GPI) anchored glycoprotein. It has a mature peptide comprising only 12 amino acids and is abundantly expressed on human lymphocytes. From the clinical point of view this protein is an important target for therapeutic interventions aimed at leukocyte depletion in hematological malignancies and post-transplant immunosuppression. CD52 / CDW52 may play a role in carrying and orienting carbohydrate. It is an unusually good target for complement-mediated cell lysis. Cancer

Immunotherapy Immune Checkpoint Immunotherapy Targeted Therapy

Reference

Piccaluga PP, et al. (2007) Expression of CD52 in peripheral T-cell lymphoma. *Haematologica*. 92(4): 566-7.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481