

Stathmin 1 Protein, Human, Recombinant (His)

General Information

Synonyms: SMN;OP18;stathmin 1;LAP18;C1orf215;PP17;Lag;PP19;PR22

Protein Construction: A DNA sequence encoding the human STMN1 (NP_005554.1) (Met1-Asp149) was expressed with a polyhistidine tag at the N-terminus. Predicted N terminal: His

Species: Human

Expression Host: *E. coli*

Accession: P16949-1

Molecular Weight: 19.5 kDa (predicted); 21-24 kDa (reducing conditions)

QC Testing

Biological Activity: Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.

Purity: > 85 % as determined by SDS-PAGE

Endotoxin: Please contact us for more information.

Formulation: Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Shipping:

In general, Lyophilized powders are shipping with blue ice.

Protein Background

Stathmin1 (STMN1) is a cytosolic phosphoprotein that regulates cellular microtubule dynamics and is known to have oncogenic activity. STMN1 is a possible biomarker for paclitaxel sensitivity and poor prognosis in GC and could be a novel therapeutic target in metastatic GC. STMN1 expression might serve as a biomarker for determining patient atypical meningioma prognosis. Stathmin1 (STMN1) is a cytosolic protein involved in microtubule dynamics through inhibition of tubulin polymerization and promotion of microtubule

depolymerization, which has been implicated in carcinogenesis and aggressive behavior in multiple epithelial malignancies. Stathmin 1 (STMN1) suppression was reported to reduce cellular viability and migration potential. STMN1 may be a promising candidate for targeted therapies in PDAC.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481