

Ezrin Protein, Human, Recombinant (His)

General Information

Synonyms: HEL-S-105;CVL;VIL2;CVIL;ezrin

Protein Construction: A DNA sequence encoding the mature form of human EZR (P15311) (Pro2-Leu586) was expressed with a polyhistidine tag at the N-terminus. Predicted N terminal: His

Species: Human

Expression Host: *E. coli*

Accession: P15311

Molecular Weight: 71.2 kDa (predicted); 82 kDa (reducing conditions)

QC Testing

Biological Activity: Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.

Purity: > 60 % as determined by SDS-PAGE

Endotoxin: Please contact us for more information.

Formulation: Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS 100 mM Arg 10% glycerol, 0.1% Tween 20, pH 7.5. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Shipping:

In general, Lyophilized powders are shipping with blue ice.

Protein Background

VIL2, also known as villin-2, belongs to the ERM protein family. It is expressed in cerebral cortex, basal ganglia, hippocampus, hypophysis, and optic nerve. VIL2 contains 1 FERM domain and may be involved in connections of major cytoskel

Reference

Grnholm M, et al. (1999) Homotypic and heterotypic interaction of the neurofibromatosis 2 tumor suppressor protein merlin and the ERM protein ezrin. *J Cell Sci.* 112(6):895-904.

Sitaraman, et al. (2002) The adenosine 2b receptor is recruited to the plasma membrane and associates with E3KARP and Ezrin upon agonist stimulation. *J Biol Chem.* 277(36):33188-95.

Gould KL, et al. (1989) cDNA cloning and sequencing of the protein-tyrosine kinase substrate, ezrin, reveals homology to band 4. *EMBO J.* 8(13):4133-42.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481