

## SLTL2/VASN Protein, Human, Recombinant (His)

### General Information

Synonyms: SLTL2;vasorin

Protein Construction: A DNA sequence encoding the human VASN (Q6EMK4) (Met1-Pro575) was expressed with a polyhistidine tag at the C-terminus. Predicted N terminal: Cys 24

Species: Human

Expression Host: HEK293 Cells

Accession: Q6EMK4

Molecular Weight: 61.1 kDa (predicted); 66-76 kDa (reducing conditions)

### QC Testing

Biological Activity: Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.

Purity: > 95 % as determined by SDS-PAGE

Endotoxin: < 1.0 EU/μg of the protein as determined by the LAL method.

Formulation: Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

### Preparation and Storage

#### Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

#### Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

#### Shipping:

In general, Lyophilized powders are shipping with blue ice.

### Protein Background

Vasorin is a typical type I membrane protein. It contains 1 EGF-like domain, 1 fibronectin type-III domain, 10 LRR (leucine-rich) repeats, 1 LRRCT domain and 1 LRRNT domain. Vasorin is predominantly expressed in vascular smooth muscle cells, and its expression is developmentally regulated. Vasorin binds to transforming growth factor (TGF)-β and attenuates TGF-β signaling in vitro. This suggests that down-regulation of vasorin expression contributes to neointimal formation after vascular injury and that vasorin modulates cellular responses.

to pathological stimuli in the vessel wall.

Reference

Gerhard DS, et al. (2004) The Status, Quality, and Expansion of the NIH Full-Length cDNA Project: The Mammalian Gene Collection (MGC) . *Genome Res.* 14(10B):2121-7.

Bauch A, et al. (2004) A physical and functional map of the human TNF-alpha/NF-kappa B signal transduction pathway. *Nat Cell Biol.* 6(2):97-105.

Nogaj LA, et al. (2005) Cellular levels of glutamyl-tRNA reductase and glutamate-1-semialdehyde aminotransferase do not control chlorophyll synthesis in *Chlamydomonas reinhardtii*. *Plant Physiol.* 139(1):389-96.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481