

## CD47 Protein, Rat, Recombinant (His)

### General Information

Synonyms:	CD47 molecule
Protein Construction:	A DNA sequence encoding the rat CD47 (P97829-1) (Met1-Lys140) was expressed, fused with a polyhistidine tag at the C-terminus. Predicted N terminal: Gln 19
Species:	Rat
Expression Host:	HEK293 Cells
Accession:	P97829-1
Molecular Weight:	15.2 kDa (predicted); 27-35 kDa (reducing condition, due to glycosylation)

### QC Testing

Biological Activity:	Measured by its binding ability in a functional ELISA. Immobilized Rat CD47-His at 10 µg/ml (100 µl/well) can bind mouse SIRPA-Fc , The EC50 of mouse SIRPA-Fc is 0.6-1.4 µg/ml.
Purity:	> 90 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

### Preparation and Storage

#### Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

#### Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

#### Shipping:

In general, Lyophilized powders are shipping with blue ice.

### Protein Background

CD47 contains 1 Ig-like V-type (immunoglobulin-like) domain and is a receptor for the C-terminal cell binding domain of thrombospondin. It may play a role in membrane transport and signal transduction. CD47 is also a membrane protein, which is involved in the increase in intracellular calcium concentration that occurs upon cell adhesion to extracellular matrix. It is very broadly distributed on normal adult tissues, as well as ovarian tumors, being especially abundant in some epithelia and the brain. CD47 may play a role in membrane transport and/or

integrin dependent signal transduction. It may prevent premature elimination of red blood cells. It also may be involved in membrane permeability changes induced following virus infection. By acting as an adhesion receptor for THBS1 on platelets, CD47 plays a role in both cell adhesion and in the modulation of integrins. It also plays an important role in memory formation and synaptic plasticity in the hippocampus. Cancer Immunotherapy Co-inhibitory Immune Checkpoint Targets Immune Checkpoint Immune Checkpoint Detection: Antibodies Immune Checkpoint Detection: ELISA Antibodies Immune Checkpoint Detection: WB Antibodies Immune Checkpoint Targets Immunotherapy Targeted Therapy

Reference

Brown EJ, et al. (2001) Integrin-associated protein (CD47) and its ligands. *Trends Cell Biol.* 11(3): 130-5.  
Oldenborg PA. (2004) Role of CD47 in erythroid cells and in autoimmunity. *Leuk Lymphoma.* 45(7): 1319-27.  
Kaczorowski DJ, et al. (2007) Targeting CD47: NO limit on therapeutic potential. *Circ Res.* 100(5): 602-3.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

**This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use**

Tel:781-999-4286 E\_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481