

## SEMA4D Protein, Rat, Recombinant (His)

### General Information

Synonyms: semaphorin 4D

Protein Construction: A DNA sequence encoding the rat SEMA4D (NP\_001164034.1)(Met1-Met712) was expressed, fused with a polyhistidine tag at the C-terminus. Predicted N terminal: Phe 24

Species: Rat

Expression Host: HEK293 Cells

Accession: A0A8I6G5Q1

Molecular Weight: 77.8 kDa (predicted); 96 kDa (reducing condition, due to glycosylation)

### QC Testing

Biological Activity: Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.

Purity: > 95 % as determined by SDS-PAGE

Endotoxin: < 1.0 EU/μg of the protein as determined by the LAL method.

Formulation: Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

### Preparation and Storage

#### Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

#### Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

#### Shipping:

In general, Lyophilized powders are shipping with blue ice.

### Protein Background

Semaphorin 4D (SEMA4D or CD100) is a member of the semaphorin family of proteins and an important mediator of the movement and differentiation of multiple cell types, including those of the immune, vascular, and nervous systems. VEGF and SEMA4D had a positive correlation with the malignant degree of ovarian cancer, and SEMA4D can serve as an independent prognostic factor. SEMA4D was the first semaphorin described to have immune functions and serves important roles in T cell priming, antibody production, and cell-to-cell adhesion. Proteolytic

cleavage of SEMA4D from the cell surface gives rise to a soluble fragment of SEMA4D (sSEMA4D). Similar to the transmembranal form, sSEMA4D is thought to have immunoregulatory properties.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481