

LILRA5/CD85f/ILT11 Protein, Rat, Recombinant (hFc)

General Information

Synonyms:	leukocyte immunoglobulin-like receptor, subfamily A (with TM domain), member 5
Protein Construction:	A DNA sequence encoding the rat LILRA5 (D4A6Y0) (Met1-Asn248) was expressed, fused with the Fc region of human IgG1 at the C-terminus. Predicted N terminal: Gln 17
Species:	Rat
Expression Host:	HEK293 Cells
Accession:	D4A6Y0
Molecular Weight:	52.8 kDa (predicted); 63 kDa (reducing conditions)

QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Shipping:

In general, Lyophilized powders are shipping with blue ice.

Protein Background

LILRA5 is a member of the leukocyte immunoglobulin-like receptor (LIR) family. LILR are a family of receptors possessing extracellular immunoglobulin domains. They are also known as CD85, ILTs, and LIR, and can exert immunomodulatory effects on a wide range of immune cells. ILT-11 contains 2 Ig-like C2-type (immunoglobulin-like) domains. It can be detected in tissues of the hematopoietic system, including bone marrow, spleen, lymph node, and peripheral leukocytes. Crosslink of ILT-11 on the surface of monocytes has been shown to induce

calcium flux and secretion of several proinflammatory cytokines, which suggests the roles of this protein in triggering innate immune responses.

Reference

Wende H, et al. (2000) Extensive gene duplications and a large inversion characterize the human leukocyte receptor cluster. *Immunogenetics*. 51(8-9):703-13.

Jones DC, et al. (2009) Alternative mRNA splicing creates transcripts encoding soluble proteins from most LILR genes. *Eur J Immunol*. 39(11):3195-206.

Mosbruger TL, et al. (2010) Large-scale candidate gene analysis of spontaneous clearance of hepatitis C virus. *J Infect Dis*. 201(9):1371-80.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481