

VEGFC Protein, Mouse/Rat, Recombinant (aa 108-223, His)

General Information

| | |
|-----------------------|--|
| Synonyms: | vascular endothelial growth factor C |
| Protein Construction: | A DNA sequence encoding the mouse / rat VEGFC (NP_033532.1/O35757) (Ala108-Arg223) was expressed fused with a polyhistidine tag at the C-terminus. Mouse and Rat mature VEGFC sequences are identical. Predicted N terminal: Ala 108 |
| Species: | Mouse,Rat |
| Expression Host: | HEK293 Cells |
| Accession: | P97953-1 |
| Molecular Weight: | 14.5 kDa (predicted) |

QC Testing

| | |
|----------------------|---|
| Biological Activity: | 1. Immobilized mouse/rat VEGFC-His at 10 µg/mL (100 µL/well) can bind mouse VEGFR3-Fc , The EC50 of mouse VEGFR3-Fc is 17.4-40.6 ng/mL. 2. Measured in a cell proliferation assay using human umbilical vein endothelial cells (HUVEC). The ED50 for this effect is typically 0.1-0.8 µg/mL. |
| Purity: | > 95 % as determined by SDS-PAGE |
| Endotoxin: | < 1.0 EU/µg of the protein as determined by the LAL method. |
| Formulation: | Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization. |

Preparation and Storage

| | |
|-----------------|---|
| Reconstitution: | A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information. |
|-----------------|---|

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Shipping:

In general, Lyophilized powders are shipping with blue ice.

Protein Background

Vascular endothelial growth factor C (VEGF-C) is a member of the VEGF family. Upon biosynthesis, VEGF-C protein is secreted as a non-covalent homodimer in an anti-parallel fashion. VEGF-C protein is a dimeric glycoprotein, as

a ligand for two receptors, VEGFR-3 (Flt4), and VEGFR-2. VEGF-C may function in angiogenesis of the venous and lymphatic vascular systems during embryogenesis. VEGF-C protein is over-expressed in various human cancers including breast cancer and prostate cancer. VEGF-C/VEGFR-3 axis, through different signaling pathways, plays a critical role in cancer progression by regulating different cellular functions, such as invasion, proliferation, and resistance to chemotherapy. Thus, targeting the VEGF-C/VEGFR-3 axis may be therapeutically significant for certain types of tumors.

Reference

- Joukov V, et al. (1997) Vascular endothelial growth factors VEGF-B and VEGF-C. *J Cell Physiol.* 173(2): 211-5.
- Su JL, et al. (2007) The role of the VEGF-C/VEGFR-3 axis in cancer progression. *Br J Cancer.* 96(4): 541-5.
- Anisimov A, et al. (2009) Activated forms of VEGF-C and VEGF-D provide improved vascular function in skeletal muscle. *Circ Res.* 104(11): 1302-12.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481