

TCCR Protein, Mouse, Recombinant (His & Avi), Biotinylated

General Information

Synonyms:	Tccr;CRL1;IL-27R;interleukin 27 receptor, α ;Wsx1;interleukin 27 receptor, alpha;zcytor1;WSX-1
Protein Construction:	A DNA sequence encoding the Mouse IL27Ra (AAC40121.1) (Met1-Lys510) was expressed with a C-terminal polyhistidine tag followed by an AVI tag. The expressed protein was biotinylated in vivo by the Biotin-Protein ligase (BirA enzyme) which is co-expressed. Predicted N terminal: Gly 29
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	AAC40121.1
Molecular Weight:	56.37 kDa (predicted); 80.69 kDa (reducing conditions)

QC Testing

Biological Activity:	Immobilized Recombinant Mouse IL27 / Interleukin-27 Protein (His Tag) at 2 μ g/mL (100 μ L/well) can bind Recombinant Mouse TCCR Protein (ECD, His & AVI Tag), Biotinylated , the EC50 is 10-30 ng/mL.
Purity:	\geq 95 % as determined by SDS-PAGE. \geq 95 % as determined by SEC-HPLC.
Endotoxin:	< 1.0 EU/ μ g of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μ m filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Shipping:

In general, Lyophilized powders are shipping with blue ice.

Protein Background

The interleukin-27 receptor is a type I cytokine receptor for interleukin-27. It is a heterodimer composed of the interleukin 27 receptor, alpha subunit, and glycoprotein 130. WSX-1/IL-27R, a class I cytokine receptor that is

homologous to the $\beta 2$ chain of the IL-12R in both sequence and structure, is highly expressed by resting/naive CD4+ T cells and CD8+ T cells. WSX-1/IL-27R belongs to the IL-6/IL-12 family of cytokines and induced proliferation of naive CD4(+) T cells and the generation of a Th1-type adaptive immune response. WSX-1/IL-27R functions as a receptor for IL27. IL-27 not only contributes to the development of an adaptive immune response through its action on CD4(+) T cells, it also directly acts on cells of the innate immune system. WSX-1/IL-27R requires IL6ST/gp130 to mediate signal transduction in response to IL27. This signaling system acts through STAT3 and STAT1. It is involved in the regulation of Th1-type immune responses. IL-27R also appears to be involved in innate defense mechanisms. IL27RA is essential for transcriptional activation of STAT1 and for augmenting the induction of T-bet expression during initiation of Th1 cell differentiation.

Reference

Pflanz S,et al.. (2004) WSX-1 and glycoprotein 130 constitute a signal-transducing receptor for IL-27. *J Immunol.* 172(4): 2225-31.

Yoshida H,et al.. (2001) WSX-1 is required for the initiation of Th1 responses and resistance to *L. major* infection. *Immunity.* 215(4): 569-78.

Villarino A,et al.. (2003) The IL-27R (WSX-1) is required to suppress T cell hyperactivity during infection. *Immunity.* 19(5): 645-55.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_email:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481